

Cohort, Exam 1

Imputed Ultrasound Data

This section contains details concerning the imputed ultrasound datasets provided on the ARIC Visit 1 data tapes. The topics covered are:

- * description of data set contents
- * data sets and variable naming conventions
- * imputed versus unimputed data

Description of Data Set Contents

Included on the updated data tapes are four data sets containing imputed ultrasound values. The data set names and variables included in each data set are described below.

Data Set and Variable Naming Conventions

Data Sets Containing Imputed Values

Because gender-race specific regression models were used to perform the imputation, a separate data set exists for White Males, White Females, Black Males, and Black Females. Each data set name consists of UBMD (indicating ultrasound) + IM (indicating the data are imputed) + WM, WF, BF, or BM (indicating the specific gender-race group). For example, the data set containing imputed ultrasound data for white males is named UBMDIMWM. Similarly, the data set containing imputed ultrasound data for black females is named UBMDIMBF. A similar pattern holds for the other gender-race groups.

The variables contained within the data sets are summarized in the table below. Most variable names consist of LBIA, RBIA, LOPA, ROPA, LINA, or RINA (indicating location) + DA or WA (indicating the type of statistic) +45 (indicating that the measurement is of the far wall). There are a few other summary variables which have unique names. These are included in the following list.

VARIABLE	DESCRIPTION	TYPE
ID	Participant ID number	Character
*DA45	Imputed site-specific average far wall thickness *=LBIA, RBIA, LOPA, ROPA, LINA, RINA	Continuous
*WA45	Weight for site-specific imputed average wall thickness *=LBIA, RBIA, LOPA, ROPA, LINA, RINA	Continuous
SUM45_1	Simple average of *DA45	Continuous
SUM45_2	Weighted average of *DA45	Continuous
SUM45_3	Z score summary statistic for *DA45	Continuous
SUMWT45	Number of observed values / 6 = weight for SUM45_1, 2, or 3	Continuous

Imputed versus Unimputed Data

You may want to rerun analyses previously run on unimputed (observed) ultrasound data (using the UBMD4 data set), on imputed data (using the UBMDIMxx data sets, where xx can be BM, BF, WM, or WF). Because of the naming conventions used, this should be a relatively easy task. Note that the data set containing unimputed ultrasound data (UBMD) contains variables of average far wall width, such as LINAAV45 and LBIAAV45. These unimputed variables on the UBMD data set correspond to the imputed variables LINADA45 and LBIADA45, respectively, on the UBMDIMxx data sets. Thus, only the middle component of the variable name must be changed for AV (unimputed average) to DA (imputed average). This logic holds true for all of the site-specific averages.

Use of Weights


The weights are a measure of precision which varies by number of sites observed. Regression estimates, using *DA45 or SUM45_1 as dependent variables, will generally be more precise if weighted regression is used.

Cohort, Exam 1**Appendix A**

B-Mode Derived Variable Site Prefixes

LAN	Left Common Carotid: Anterior Angle
RAN	Right Common Carotid: Anterior Angle
LBI	Left Bifurcation
RBI	Right Bifurcation
LIN	Left Internal Carotid
RIN	Right Internal Carotid
LOP	Left Common Carotid: Optimal Angle
ROP	Right Common Carotid: Optimal Angle
LPO	Left Common Carotid: Posterior Angle
RPO	Right Common Carotid: Posterior Angle
LPP	Left Popliteal
RPP	Right Popliteal
QC1	First QC Repeat Scan (refer to QC01 for site identification)
QC2	Second QC Repeat Scan (refer to QC02 for site identification)

Schematic Overview of Carotid Artery B-Mode Ultrasound Measurements



- Interfaces:
- 1- Boundary between the periadventitia and adventitia of the near wall (not measured)
 - 2- Boundary between the adventitia and media of the near wall
 - 3- Boundary between the intima of the near wall and the blood
 - 4- Boundary between blood and intima of the far wall
 - 5- Boundary between media and adventitia of the far wall
 - 6- Boundary between adventitia and periadventitia of the far wall (not measured)

Max 23 = B-A; Max 45 = D-C; Min 34 = H-G

The extracranial carotid system is divided into one-centimeter segments: I = internal carotid; II = carotid bifurcation; III = common carotid. A maximum of eleven measurements is made by URC readers on each arterial wall interface, in each arterial segment. These measurements are placed equidistant at 1 millimeter intervals, represented by the eleven points placed on interface B2 on the internal carotid. Also shown on this schematic is the definition of a maximum and a minimum wall thickness variable.

Cohort, Exam 1**Ultrasound data**

Imputed, white male

ID		Aric Subject ID (Cir)
N	Value	Description
5330	Present	Text suppressed

LBIADA45		Derived Average Far Wall Thickness, Left Bifurcation
N	Value	Description
5112	Range	0.2345 - 4.12529 (median=0.848667 mean=0.9054743 std=0.3372658)
218		Missing

LBIAWA45		Weight For LBIADA45
N	Value	Description
380	0.1666666667	
528	0.3333333333	
546	0.5	
414	0.6666666667	
151	0.8333333333	
3093	1	
218		Missing

LINADA45		Derived Average Far Wall Thickness, Left Internal Carotid
N	Value	Description
5112	Range	0.201 - 3.79667 (median=0.689231 mean=0.7425888 std=0.2883006)
218		Missing

LINAWA45		Weight For LINADA45
N	Value	Description
402	0.1666666667	
642	0.3333333333	
742	0.5	
671	0.6666666667	
351	0.8333333333	
2304	1	
218		Missing

Cohort, Exam 1

<i>LOPADA45</i>		
<i>N</i>	<i>Value</i>	<i>Description</i>
5112	Range	0.268 - 2.09527 (median=0.682247 mean=0.7052656 std=0.1751984)
218		Missing

<i>LOPAWA45</i>		
<i>N</i>	<i>Value</i>	<i>Description</i>
268	0.1666666667	
293	0.3333333333	
208	0.5	
91	0.6666666667	
47	0.8333333333	
4205	1	
218		Missing

<i>RBIADA45</i>		
<i>N</i>	<i>Value</i>	<i>Description</i>
5112	Range	0.156333 - 4.70827 (median=0.866297 mean=0.9384666 std=0.3823429)
218		Missing

<i>RBIAWA45</i>		
<i>N</i>	<i>Value</i>	<i>Description</i>
381	0.1666666667	
551	0.3333333333	
512	0.5	
330	0.6666666667	
139	0.8333333333	
3199	1	
218		Missing

Cohort, Exam 1

RESPONS		<i>Number Of Observed Sites</i>
N	Value	Description
218	0	
421	1	
736	2	
993	3	
1146	4	
1080	5	
736	6	

RINADA45		<i>Derived Average Far Wall Thickness, Right Internal Carotid</i>
N	Value	Description
5112	Range	0.1675 - 5.695 (median=0.757881 mean=0.8241107 std=0.3582125)
218		Missing

RINAWA45		<i>Weight For RINADA45</i>
N	Value	Description
400	0.1666666667	
655	0.3333333333	
743	0.5	
650	0.6666666667	
332	0.8333333333	
2332	1	
218		Missing

ROPADA45		<i>Derived Average Far Wall Thickness, Right Common Carotid: Optimal Angle</i>
N	Value	Description
5112	Range	0.2345 - 3.551 (median=0.658349 mean=0.6805581 std=0.1757669)
218		Missing

Cohort, Exam 1

ROPAWA45		Weight For ROPADA45
N	Value	Description
274	0.1666666667	
275	0.3333333333	
228	0.5	
136	0.6666666667	
60	0.8333333333	
4139	1	
218		Missing

SUM45_1		Mean Of The DA45 Variables
N	Value	Description
5112	Range	0.399947 - 2.812405 (median=0.754865 mean=0.7994107 std=0.2073423)
218		Missing

SUM45_2		Weighted Mean Of The DA45 Variables
N	Value	Description
5112	Range	0.403196 - 2.855114 (median=0.756752 mean=0.7995014 std=0.2022818)
218		Missing

SUM45_3		Z-Score Summary Statistic For The DA45 Variables
N	Value	Description
5112	Range	0.359667 - 3.089784 (median=0.756817 mean=0.7994626 std=0.2127214)
218		Missing

SUMWT45		'Number of observed values / 6 = weight for Sum45_1
N	Value	Description
421	0.1666666667	
736	0.3333333333	
993	0.5	
1146	0.6666666667	
1080	0.8333333333	
736	1	
218		Missing