

ARIC Manuscript Proposal #1919

PC Reviewed: 3/20/2012
SC Reviewed: _____

Status: A
Status: _____

Priority: 2
Priority: _____

1.a. Full Title:

Prognostic Significance of Bundle Branch Blocks as Independent Predictors of Incident Heart Failure in the Atherosclerosis Risk in Communities Study (ARIC)

b. Abbreviated Title (Length 26 characters):

Bundle Blocks & Heart failure

2. Writing Group:

Writing group members:

Zhu-ming Zhang, Pentti Rautaharju, Ronald J Prineas, Laura Loehr,
Wayne Rosamond, Elsayed Z Soliman

I, the first author, confirm that all the coauthors have given their approval for this manuscript proposal. ZMZ [please confirm with your initials electronically or in writing]

First author:

Zhu-Ming Zhang, MD, MPH

Address:

Assistant Professor
Epidemiological Cardiology Research Center (EPICARE)
Department of Epidemiology and Prevention
Division of Public Health Sciences
Wake Forest School of Medicine
Medical Center Blvd. (Wells Fargo-13)
Winston-Salem, NC 27157
Phone: (336)716-0835 Fax: (336) 716-0834
Email: zmzhang@wakehealth.edu

Corresponding author (must be an ARIC investigator for the proposal but can be different in the published paper; correspondence will be sent to both the first author & the corresponding author):

Elsayed Z Soliman, MD, MSc, MS
Epidemiological Cardiology Research Center (EPICARE)
Wake Forest School of Medicine
Medical Center Blvd. (Well Fargo-13)
Winston-Salem, NC 27157
Phone: (336)716-8632 Fax: (336) 716-0834

3. Timeline:

Start analyses: upon receipt of data from the coordinating centre
Submission for publication: December 2012

4. Rationale:

Cardiovascular disease continues to be a major cause of death and disability worldwide, with a rising epidemic of heart Failure (HF) in aging populations¹ Most HF is a consequence of ischemic, or hypertensive. Abnormalities of the ST-T waves, and widening of the QRS/T angle in the resting electrocardiogram (ECG) have been associated with incident heart failure, coronary heart disease (CHD), and all-cause mortality in the general population.²⁻⁷ Furthermore, bundle branch blocks (BBBs), both left (LBBB) and right (RBBB), have been shown to be predictive of all-cause mortality and incident HF in selected populations⁸⁻¹⁷

However, the presence of complete BBB is thought to limit the utility of the ECG in evaluating myocardial ischemia, where it is more difficult to categorize the accompanying appearance of abnormal depolarization and repolarization of the heart. LBBB, in particular, is characterized by repolarization changes in which the ST segments are discordant to the major direction of the QRS complex. Although the prognostic significance of these repolarization abnormalities associated with BBB have been shown for mortality and overall CHD in postmenopausal women⁹, these abnormalities have not been specifically tested for measuring ischemia as a predictor of CHF in men or women. Also it is not entirely clear whether different patterns of BBB (LBBB, RBBB, intraventricular conduction defects (IVCD)) or prevalent versus incident BBB have different predictive value or not.

The aim of this study is to evaluate the prognostic significance of prevalent and incident BBB (LBBB, RBBB and IVCD, separately) in predicting incident HF, and to evaluate whether the prognostic value of BBB for HF will be enhanced by combining with ischemia-related ECG parameters including QRS/T angle (spatial or frontal planes), discordance of repolarization by various measures such as maximal ST and T deviation and the net QRS integral in representative leads of various lead groups, and rate-adjusted QT interval, separately.

5. Main Hypothesis/Study Questions:

This study aims to:

- (1) Estimate the prognostic significance of baseline '*prevalent*' LBBB, RBBB, and IVCD, separately, for the prediction of incident heart failure in the ARIC study.
- (2) Estimate the prognostic significance of '*incident*' LBBB, RBBB, and IVCD, separately, for the prediction of incident heart failure in the ARIC study.
- (3) Examine the added predictive value of combining BBB (LBBB, RBBB, and IVCD, separately) with repolarization abnormalities (spatial and frontal QRS/T angle maximal ST and T deviation and the net QRS integral in representative leads of various lead groups, and rate-adjusted QT interval) for prediction of heart failure in the ARIC study.

6. Design and analysis (study design, inclusion/exclusion, outcome and other variables of interest with specific reference to the time of their collection, summary of data analysis, and any anticipated methodological limitations or challenges if present).

Sample Size

All ARIC participants with good quality baseline ECG data as well as information on all relevant risk factors and outcome events of CHF during ARIC follow-up will be eligible for inclusion in this analysis. Participants with artificial pacemaker, or Wolf Parkinson White Syndrome will be excluded. From the total of 15,571 ARIC participants at baseline we expected to have approximately 450 ECGs with BBB, and 2000 ECGs with QRS duration ≥ 120 ms through 4 visits. There are approximately 1800 incident heart failures in the total cohort during a maximum of 17 years follow-up through 2006.

The Variables:

ECG variables needed to fulfill the aim of the study will be:

Complete Minnesota codes¹⁸⁻¹⁹ for all ECGs at baseline and follow up visits, which will include specifically,

- Complete left bundle branch block (MC-7.1)
- Complete right bundle branch block (MC-7.2)
- Combination of RBBB and left anterior fascicular block (MC-7.8)
- Intraventricular conduction defect, QRS ≥ 120 ms (MC-7.4).
- ECG- Myocardial infarction/Ischemia -- MC 1, MC 4, MC 5, MC 92 --Q/QS wave, ST segment, T wave amplitude.
- Frontal QRS axis and T axis, and QRS/T angles (spatial and frontal)
- Continuous measurements of the duration and amplitude of Q-R-S-T wave by lead.
- Presence of arrhythmias and ectopic beat including atrial and ventricular ectopic beats and atrial fibrillation.

Non-ECG variables:

Non-ECG variables include demographic and clinic data, and outcome measures, which are summarized in below:

- (1) The key demographic and clinical variables -- age, race, gender, body mass index, education, smoking status, alcohol use, hypertension, diabetes mellitus, cancer, emphysema, previous stroke, history of cardiovascular disease, family history of coronary heart disease, HDL cholesterol, LDL cholesterol, total triglycerides, total cholesterol, systolic blood pressure, diastolic blood pressure, hematocrit, white blood cell, baseline fasting blood glucose, insulin, creatinine, fibrinogen, and uric acid.
- (2) The outcome includes updated incident heart failure confirmed by the endpoints committee (update to the end of 2006).

INCHF06	Incident HF (from discharge codes)
---------	------------------------------------

Data analysis:

First, frequency distributions of all ECG and non-ECG variables will be inspected to rule out anomalies and outliers possibly due to measurement artifacts.

Time dependent Cox regression analysis will be used to examine the risk of incident heart failure and the association with each pattern of BBB (LBBB, RBBB, and IVCD) as predictors for incident heart failure.

Models will be initially adjusted for demographic (age, sex, race), then further adjusted for clinical characteristics (which are the variables mentioned as non-ECG variables above). Similar analysis will be conducted for QRS duration, QRS/T angle (spatial and frontal), amplitude of Q-R-S by leads, magnitude of ST segment elevation or depression in leads with predominantly positive or negative QRS complexes (discordant or concordant of QRS complex and ST segment), magnitude of T wave change in leads with predominantly positive or negative QRS complexes (discordant or concordant of QRS complex and T wave), separately and combined with BBB patterns. Interaction by sex, race, and history of CVD will be examined and models will be stratified accordingly.

The proportional hazards assumption of the Cox regression model will be checked graphically for each of the candidate variables. All analyses will be performed with the SAS software, version 9.2.

7.a. Will the data be used for non-CVD analysis in this manuscript? ___ Yes ☒ No

b. If Yes, is the author aware that the file ICTDER03 must be used to exclude persons with a value RES_OTH = "CVD Research" for non-DNA analysis, and for DNA analysis RES_DNA = "CVD Research" would be used?

___ Yes ___ No

(This file ICTDER03 has been distributed to ARIC PIs, and contains the responses to consent updates related to stored sample use for research.)

8.a. Will the DNA data be used in this manuscript? ___ Yes ☒ No

8.b. If yes, is the author aware that either DNA data distributed by the Coordinating Center must be used, or the file ICTDER03 must be used to exclude those with value RES_DNA = "No use/storage DNA"?

___ Yes ___ No

9. The lead author of this manuscript proposal has reviewed the list of existing ARIC Study manuscript proposals and has found no overlap between this proposal and previously approved manuscript proposals either published or still in active status.

ARIC Investigators have access to the publications lists under the Study Members Area of the web site at: <http://www.csc.c.unc.edu/ARIC/search.php>

☒ Yes ___ No

10. What are the most related manuscript proposals in ARIC (authors are encouraged to contact lead authors of these proposals for comments on the new proposal or collaboration)?

11. a. Is this manuscript proposal associated with any ARIC ancillary studies or use any ancillary study data? _____ Yes ☒ No

11.b. If yes, is the proposal

_____ **A. primarily the result of an ancillary study (list number* _____)**

_____ **B. primarily based on ARIC data with ancillary data playing a minor role (usually control variables; list number(s)* _____)**

*ancillary studies are listed by number at <http://www.csc.unc.edu/aric/forms/>

12. Manuscript preparation is expected to be completed in one to three years. If a manuscript is not submitted for ARIC review at the end of the 3-years from the date of the approval, the manuscript proposal will expire.

References:

- 1- National Institutes of Health, National Heart, Lung and Blood Institute, Morbidity and Mortality: 2009 Chart book on Cardiovascular, Lung and Blood Diseases. Bethesda MD: National Institutes of Health: 2009
- 2- Rautaharju, PM, Kooperberg C, Larson JC, LaCroix, A. Electrocardiographic Predictors of Incident Congestive Heart Failure and All-cause Mortality in Postmenopausal Women. The Women's Health Initiative. *Circulation* 2006;113:481-489.
- 3- Rautaharju PM, Prineas RJ, Wood J, et al. Electrocardiographic predictors of new-onset heart failure in men and women free of coronary heart disease (from the Atherosclerosis in Communities [ARIC] Study). *Am J Cardiol* 2007;100:1437-1441.
- 4- Zhang ZM, Prineas RJ, Case D, Soliman EZ, Rautaharju PM for the ARIC Research Group: Comparison of the prognostic significance of the electrocardiographic QRS/Tangles in predicting incident coronary heart disease and total mortality (from the atherosclerosis risk in communities study). *Am J Cardiol* 2007;100:844-849
- 5- Zhang ZM, Prineas RJ, Eaton CB: Evaluation and comparison of the Minnesota code and Novacode for electrocardiographic Q-ST wave abnormalities for the independent prediction of incident coronary heart disease and total mortality (from the Women's Health Initiative). *Am J Cardiol* 2010;106(1):18-25.
- 6- Kors JA, van Herpen G, Sittig AC et al., Reconstruction of the Frank vectorcardiogram from standard electrocardiographic leads: diagnostic comparison of different methods, *Eur Heart J* 1990:1083.
- 7- Schreurs CA, Algra AM, Man SC, Cannegieter SC, van der Wall EE, Schalij MJ, Kors, JA, Swenne CA. The spatial QRS-T angle in the Frank vectorcardiogram: accuracy of estimates derived from the 12-lead electrocardiogram. *J Electrocardiol* 2010;43(4):294-301.

- 8- Pfister R, Cairns R, Erdmann E, Schneider CA. A clinical risk score for heart failure in patients with type 2 diabetes and macrovascular disease: An analysis of the PROactive study. *Int J Cardiol.* 2011 May 3
- 9- Zhang ZM, Rautaharju PM, Soliman EZ, Manson JE, Cain ME, Martin LW, Bavry A, Metha L, Vitolins M, Prineas RJ. Mortality Risk Associated with Bundle Branch Blocks and Associated Repolarization Abnormalities: The Women's Health Initiative (WHI). (under journal review)
- 10- Baldasseroni S, De Biase L, Fresco C, Marchionni N, Marini M, Masotti G, Orsini G, Porcu M, Pozzar F, Scherillo M, Maggioni AP; Italian Network on Congestive Heart Failure. Cumulative effect of complete left bundle-branch block and chronic atrial fibrillation on 1-year mortality and hospitalization in patients with congestive heart failure. A report from the Italian network on congestive heart failure (in-CHF database). *Eur Heart J.* 2002;23:1692–1698.
- 11- Zannad F, Huvelle E, Dickstein K, van Veldhuisen DJ, Stellbrink C, Køber L, Cazeau S, Ritter P, Maggioni AP, Ferrari R, Lechat P. Left bundle branch block as a risk factor for progression to heart failure. *European Journal of Heart Failure.* 2007;9:7-14
- 12- Tabrizi F, Englund A, Rosenqvist M, Wallentin L, Stenestrand U: Influence of left bundle branch block on long-term mortality in a population with heart failure. *Eur Heart J.* 2007; 28 (20): 2449-2455.
- 13- Clark AL, Goode K, Cleland JGF. The prevalence and incidence of left bundle branch block in ambulant patients with chronic heart failure. *Eur J Heart Fail.* 2008;10(7):696-702.
- 14- Imanishi R, Seto S, Ichimaru S, Nakashima E, Yano K, Akahoshi M. Prognostic significance of incident complete left bundle branch block observed over a 40-year period. *Am J Cardiol.* 2006;98:644.
- 15- Agarwal SK, Simpson RJ Jr, Rautaharju P, Alonso A, Shahar E, Massing M, Saba S, Heiss G. Relation of Ventricular Premature Complexes to Heart Failure (from the Atherosclerosis Risk In Communities [ARIC] Study). *Am J Cardiol.* 2011 Sep 24.
- 16- Sosin M, Davis RC, Lip GYH. Mortality in congestive heart failure complicated by atrial fibrillation. *Eur Heart J.* 2004;25(1):97-98
- 17- Auricchio A, Fantoni C, Regoli F, et al. Characterization of left ventricular activation in patients with heart failure and left bundle-branch block. *Circulation* 2004; 109:1133.
- 18- Prineas RJ, Crow RS, Blackburn H. The Minnesota Code Manual of Electrocardiographic Findings. Boston: John Wright PSB, 1982;203
- 19- Prineas RJ, Crow RS, Zhang ZM. The Minnesota Code Manual of Electrocardiographic Findings (Second edition). Published by Springer-London, 2009.